

**STUDY OF A COLLECTION OF SERVICE TIME DISTRIBUTIONS AND
 IMPACT IN THE RESPECTIVE M|G| ∞ SYSTEM BUSY PERIOD
 PARAMETERS**

FERREIRA Manuel Alberto M. (PT), FILIPE José António (PT)

Abstract. The problems arising when computing the moments of a particular service time distributions collection, for which the M|G| ∞ queue system busy period becomes very easy to study, are presented and it is shown how to overcome them. Some results, precisely about the moment's computation of random variables with distribution functions given by this collection are given. The busy period “peakedness” and “modified peakedness” for the M|G| ∞ queue in the case of those service time distributions are also computed.

Keywords: service time, collection, distributions, moments, M|G| ∞ queue, busy period

Mathematics Subject Classification: 60K35

1 Introduction

If, in the M|G| ∞ queue system, the service time length is a random variable with a distribution function belonging to the collection

$$G(t) = 1 - \frac{(1 - e^{-\rho}) \left(\lambda + \frac{\lambda p + \beta}{1 - p} \right)}{\lambda e^{-\rho} \left(e^{\left(\lambda + \frac{\lambda p + \beta}{1 - p} \right) t} - 1 \right) + \lambda}, t \geq 0, -\lambda \leq \beta \leq \frac{\lambda(1 - pe^\rho)}{e^\rho - 1}, 0 \leq p < 1 \quad (1.1),$$

the busy period length probability distribution is exponential with an atom at the origin, see Ferreira (2005) and (Ferreira and Andrade, 2009). But although it is so easy to study the busy period in this situation it is very difficult to compute the service time moments.

Some results, precisely about the moment's computation of random variables with distribution functions given by this collection are given.

In the end are presented formulae that give the “peakedness” and the “modified peakedness” to the busy period of the $M|G|^\infty$ system for those service time distributions, see Ferreira (2004, 2013,2013a).

This work is built on a part of the presented in Ferreira (2007) which is so corrected, generalized and updated.

2 Moments Calculation Problems

Be $G(t), t \geq 0$ a distribution function and $g(t) = \frac{dG(t)}{dt}$. The differential equation

$$(1-p) \frac{g(t)}{1-G(t)} - \lambda p - \lambda(1-p)G(t) = \beta, \text{ where } \lambda > 0 \text{ and } -\lambda \leq \beta \leq \frac{\lambda(1-pe^\rho)}{e^\rho - 1}, \quad 0 \leq p < 1$$

($\rho = \lambda\alpha$, being α the mean of $G(t)$) has (1.1) as solution (see Ferreira (2005)). If, in (1.1), $G_i(t)$ is the solution associated to ρ_i , $i = 1,2,3,4$ it is easy to see that

$$\frac{G_4(t) - G_2(t)}{G_4(t) - G_1(t)} \cdot \frac{G_3(t) - G_1(t)}{G_3(t) - G_2(t)} = \frac{e^{-\rho_4} - e^{-\rho_2}}{e^{-\rho_4} - e^{-\rho_1}} \cdot \frac{e^{-\rho_3} - e^{-\rho_1}}{e^{-\rho_3} - e^{-\rho_2}} \quad (2.1)$$

as it had to happen since the differential equation dealt with is a Riccati equation. And computing,

$$\begin{aligned} \int_0^\infty [1 - G(t)] dt &= \int_0^\infty \frac{(1 - e^{-\rho}) \left(\lambda + \frac{\lambda p + \beta}{1 - p} \right)}{\lambda e^{-\rho} \left(e^{\left(\lambda + \frac{\lambda p + \beta}{1 - p} \right) t} - 1 \right) + \lambda} dt = \\ &= \frac{(1 - e^{-\rho}) \left(\lambda + \frac{\lambda p + \beta}{1 - p} \right)}{\lambda} \int_0^\infty \frac{1}{e^{-\rho} \left(e^{\left(\lambda + \frac{\lambda p + \beta}{1 - p} \right) t} - 1 \right) + 1} dt = \\ &= \frac{(1 - e^{-\rho}) \left(\lambda + \frac{\lambda p + \beta}{1 - p} \right)}{\lambda} \int_0^\infty \frac{e^{-\left(\lambda + \frac{\lambda p + \beta}{1 - p} \right) t}}{e^{-\rho} - e^{-\rho} e^{-\left(\lambda + \frac{\lambda p + \beta}{1 - p} \right) t} + e^{-\left(\lambda + \frac{\lambda p + \beta}{1 - p} \right) t}} dt = \end{aligned}$$

$$\begin{aligned}
&= \frac{(1-e^{-\rho})\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)}{\lambda} \int_0^{\infty} \frac{e^{-\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)t}}{e^{-\rho} + (1-e^{-\rho})e^{-\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)t}} dt = \\
&= \frac{(1-e^{-\rho})\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)}{\lambda} \cdot \frac{-1}{(1-e^{-\rho})\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)} \cdot \left[\log \left(e^{-\rho} + (1-e^{-\rho})e^{-\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)t} \right) \right]_0^{\infty} = \\
&= -\frac{1}{\lambda} (\log e^{-\rho} - \log 1) = \frac{-\rho}{-\lambda} = \alpha
\end{aligned}$$

as it had to be because are considered positive random variable. The density associated to $G(t)$ given by (1.1) is

$$g(t) = \frac{(1-e^{-\rho})e^{-\rho}\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)^2 e^{-\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)t}}{\lambda \left[e^{-\rho} + (1-e^{-\rho})e^{-\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)t} \right]^2}, t > 0, -\lambda \leq \beta \leq \frac{\lambda(1-pe^{-\rho})}{e^{\rho}-1}, 0 \leq p < 1 \quad (2.2).$$

So,

$$\int_0^{\infty} t^n g(t) dt = \frac{(1-e^{-\rho})e^{-\rho}\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)^2}{\lambda} \cdot \int_0^{\infty} t^n \frac{e^{-\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)t}}{\left[e^{-\rho} + (1-e^{-\rho})e^{-\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)t} \right]^2} dt.$$

$$\text{But, } \int_0^{\infty} t^n \frac{e^{-\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)t}}{\left[e^{-\rho} + (1-e^{-\rho})e^{-\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)t} \right]^2} dt \geq \int_0^{\infty} t^n e^{-\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)t} dt =$$

$$= \frac{1}{\lambda + \frac{\lambda p + \beta}{1-p}} \frac{n!}{\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)^n}, \beta \neq -\lambda. \text{ And,}$$

$$\int_0^{\infty} t^n \frac{e^{-\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)t}}{\left[e^{-\rho} + (1 - e^{-\rho}) e^{-\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)t} \right]^2} dt \leq e^{2\rho} \int_0^{\infty} t^n e^{-\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)t} dt =$$

$$= \frac{e^{2\rho}}{\lambda + \frac{\lambda p + \beta}{1-p}} \frac{n!}{\left(\lambda + \frac{\lambda p + \beta}{1-p} \right)^n}, \beta \neq -\lambda.$$

So, calling T the random variable corresponding to $G(t)$:

$$\frac{(1 - e^{-\rho}) e^{-\rho}}{\lambda} \frac{n!}{\left(\lambda + \frac{\lambda p + \beta}{1-p} \right)^{n-1}} \leq E[T^n] \leq \frac{e^{\rho} - 1}{\lambda} \frac{n!}{\left(\lambda + \frac{\lambda p + \beta}{1-p} \right)^{n-1}},$$

$$, -\lambda < \beta \leq \frac{\lambda(1 - pe^{-\rho})}{e^{\rho} - 1}, 0 \leq p < 1, n = 1, 2, \dots \quad (2.3).$$

Notes:

- The expression (2.3), giving bounds for $E[T^n]$, guarantees its existence,
- For $n = 1$ the expression (2.3) is useless since $E[T] = \alpha$. Note, curiously, that here the upper bound is $\frac{e^{\rho} - 1}{\lambda}$, the M|G| ∞ system busy period mean value, evidently disparate,
- For $n = 2$, subtracting to both bounds α^2 , it is possible get from expression (2.3) bounds for $VAR[T]$,
- For $\beta = -\lambda, E[T^n] = 0, n = 1, 2, \dots$, evidently.

See, however, that (1.1) can be written like:

$$G(t) = \frac{1 + \frac{\lambda p + \beta}{1-p} (1 - e^{\rho}) e^{-\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)t}}{1 - (1 - e^{\rho}) e^{-\left(\lambda + \frac{\lambda p + \beta}{1-p}\right)t}}, t \geq 0, -\lambda \leq \beta \leq \frac{\lambda(1 - pe^{\rho})}{e^{\rho} - 1}, 0 \leq p < 1 \quad (2.4)$$

and, for $\rho < \log 2$,

$$G(t) = \left(1 + \frac{\lambda p + \beta}{\lambda} (1 - e^\rho) e^{-\left(\lambda + \frac{\lambda p + \beta}{1 - p}\right)t} \right) \cdot \sum_{k=0}^{\infty} (1 - e^\rho)^k e^{-k\left(\lambda + \frac{\lambda p + \beta}{1 - p}\right)t},$$

$$, t \geq 0, -\lambda \leq \beta \leq \frac{\lambda(1 - pe^\rho)}{e^\rho - 1}, 0 \leq p < 1 \quad (2.5).$$

After (2.5) it is easy to derive the T Laplace Transform for $\rho < \log 2$. And, so,

- For $\rho < \log 2$

$$E[T^n] = - \left(1 + \frac{\lambda p + \beta}{\lambda} \right) n! \sum_{k=1}^{\infty} \frac{(1 - e^\rho)^k}{k \left(\lambda + \frac{\lambda p + \beta}{1 - p} \right)^n}, -\lambda < \beta \leq \frac{\lambda(1 - pe^\rho)}{e^\rho - 1}, 0 \leq p < 1,$$

$$, n = 1, 2, \dots \quad (2.6).$$

Notes:

$$- E[T] = - \left(1 + \frac{\lambda p + \beta}{\lambda} \right) \sum_{k=1}^{\infty} \frac{(1 - e^\rho)^k}{k \left(\lambda + \frac{\lambda p + \beta}{1 - p} \right)} = \frac{1}{\lambda} \sum_{k=1}^{\infty} (-1)^{k+1} \frac{(1 - e^\rho)^k}{k} =$$

$$= \frac{1}{\lambda} \log e^\rho = \frac{\rho}{\lambda} = \alpha.$$

- For $n \geq 2$ only a finite number of parcels can be considered in the infinite sum. Calling M this number, to get an error lesser than ε it must be fulfilled simultaneously

$$a) M > \frac{1}{\lambda + \frac{\lambda p + \beta}{1 - p}} - 1,$$

$$b) M > \log_{(e^\rho - 1)} \frac{\varepsilon e^\rho \lambda}{n! \left(\lambda + \frac{\lambda p + \beta}{1 - p} \right)} - 1.$$

So it is evident now that this distributions collection moment's computation is a complex task. This was already true for the study of Ferreira (1998) where the results presented are a particular situation of these ones for $p = 0$. To consider the approximation

$$E_m^n = \sum_{k=1}^{\infty} \left(\frac{k}{m}\right)^n \left[G\left(\frac{k}{m}\right) - G\left(\frac{k-1}{m}\right) \right], -\lambda < \beta \leq \frac{\lambda(1-pe^\rho)}{e^\rho-1}, 0 \leq p < 1, n = 1, 2, \dots \quad (2.7)$$

may be helpful since $\lim_{m \rightarrow \infty} E_m^n = E[T^n]$, $n = 1, 2, \dots$ (Ferreira and Andrade, 2012c) that allow the moments numerical computation .

3 The “Peakedness” and the “Modified Peakedness” for the M|G| ∞ Queue Busy Period

The M|G| ∞ queue busy period “peakedness” is the Laplace transform of its length at $\frac{1}{\alpha}$, Ferreira (2013,2013a) . It is a parameter that characterizes the busy period distribution length and contains information about all its moments. For the collection of service distributions (1.1) the “peakedness”, named pi , is

$$pi = \frac{e^{-\rho}(\lambda + \beta)(\rho + 1) - \lambda p - \beta}{\lambda(e^{-\rho}(\rho + \alpha\beta) + 1 - p)}, -\lambda \leq \beta \leq \frac{\lambda(1-pe^\rho)}{e^\rho-1}, 0 \leq p < 1 \quad (3.1).$$

In Ferreira (2013,2013a) is also introduced another measure, the “modified peakedness” got after the “peakedness” taking out the terms that are permanent for the busy period in different service distributions and putting over the common part. Calling it qi :

$$qi = pi \frac{\rho}{e^\rho - \rho - 1} + 1$$

and so, for the distributions given by collection (1.1):

$$qi = \frac{e^{-\rho}(\lambda + \beta)(\rho + 1) - \lambda p + \beta}{\lambda(e^{-\rho}(\rho + \alpha\beta) + 1 - p)} \frac{\rho}{e^\rho - \rho - 1} + 1, -\lambda \leq \beta \leq \frac{\lambda(1-pe^\rho)}{e^\rho-1}, 0 \leq p < 1 \quad (3.2).$$

References

- [1] ANDRADE, M. (2010). A note on foundations of probability. *Journal of Mathematics and Technology*, 1(1), 96-98.
- [2] CARRILLO, M. J. (1991). Extensions of Palm’s theorem: a review. *Management Science*, 37(6), 739-744.

- [3] FERREIRA, M. A. M., RAMALHOTO, M. F. (1994). Estudo dos parâmetros básicos do período de ocupação da fila de espera $M|G|\infty$. *A Estatística e o Futuro e o Futuro da Estatística*. Actas do I Congresso Anual da S.P.E., Edições Salamandra, Lisboa, 47-60.
- [4] FERREIRA, M. A. M. (1998). Momentos de Variáveis Aleatórias com Função de Distribuição dadas pela Coleção $G(t) = 1 - \frac{(1 - e^{-\rho})(\lambda + \beta)}{\lambda e^{-\rho}(e^{(\lambda + \beta)} - 1) + \lambda}, t \geq 0, -\lambda \leq \beta \leq \frac{\lambda}{e^{\rho} - 1}$. *Revista Portuguesa de Gestão*, II. ISCTE, Lisboa, 67-69.
- [5] FERREIRA, M. A. M. (1998a). Application of Riccati equation to the busy period study of the $M|G|\infty$ system. *Statistical Review*, 1st Quadrimester, INE, 23-28.
- [6] FERREIRA, M. A. M. (1998b). Computational simulation of infinite servers systems. *Statistical Review*, 3rd Quadrimester, INE, 23-28.
- [7] FERREIRA, M. A. M. (2001). $M|G|\infty$ queue heavy-traffic situation busy period length distribution (power and Pareto service distributions). *Statistical Review*, 1st Quadrimester, INE, 27-36.
- [8] FERREIRA, M. A. M. (2002). The exponentiality of the $M|G|\infty$ queue busy period. *Actas das XII Jornadas Luso-Espanholas de Gestão Científica*, Volume VIII-Economia da Empresa e Matemática Aplicada. UBI, Covilhã, Portugal, 267-272.
- [9] FERREIRA, M. A. M. (2004). $M|G|\infty$ Queue Busy Cycle Renewal Function for Some Particular Service Time Distributions. *Proceedings of Quantitative Methods in Economics (Multiple Criteria Decision Making XII)*. Virt, Slovakia, 42-47.
- [10] FERREIRA, M. A. M. (2005). Differential equations important in the $M|G|\infty$ queue system transient behavior and busy period study. *Proceedings of 4th International Conference APLIMAT 2005*, Bratislava, Slovakia, 119-132.
- [11] FERREIRA, M. A. M. (2005a). A Equação de Riccati no Estudo da Ocupação de um Sistema $M|G|\infty$ (Uma Generalização). *Actas do I Congresso de Estatística e Investigação Operacional da Galiza e Norte de Portugal/VII Congresso Galego de Estatística e Investigación de Operacións*. Guimarães, Portugal.
- [12] FERREIRA, M. A. M. (2007). $M|G|\infty$ system parameters for a particular collection of service time distributions. *Proceedings of 6th International Conference APLIMAT 2007*, Bratislava, Slovakia, 131-137.
- [13] FERREIRA, M. A. M., ANDRADE, M., FILIPE, J. A. (2009). Networks of queues with infinite servers in each node applied to the management of a two echelons repair system. *China-USA Business Review*, 8(8), 39-45 and 62.
- [14] FERREIRA, M. A. M., ANDRADE, M. (2009). The ties between the $M|G|\infty$ queue system transient behavior and the busy period. *International Journal of Academic Research*, 1(1), 84-92.
- [15] FERREIRA, M. A. M., ANDRADE, M. (2010). Looking to a $M|G|\infty$ system occupation through a Riccati equation. *Journal of Mathematics and Technology*, 1 (2), 58-62.

- [16] FERREIRA, M. A. M., ANDRADE, M. (2010a). $M|G|\infty$ system transient behavior with time origin at the beginning of a busy period mean and variance. *APLIMAT-Journal of Applied Mathematics*, 3(3), 213-221.
- [17] FERREIRA, M. A. M., ANDRADE, M. (2011). Fundamentals of theory of queues. *International Journal of Academic Research*, 3(1), part II, 427-429.
- [18] FERREIRA, M. A. M., ANDRADE, M. (2012). Busy period and busy cycle distributions and parameters for a particular $M|G|\infty$ queue system. *American Journal of Mathematics and Statistics*, 2(2), 10-15. <http://article.sapub.org/10.5923.j.ajms.20120202.03.html>
- [19] FERREIRA, M. A. M., ANDRADE, M. (2012a). Queue networks models with more general arrival rates. *International Journal of Academic Research*, 4(1), part A, 5-11.
- [20] FERREIRA, M. A. M., ANDRADE, M. (2012b). Transient behavior of the $M|G|m$ and $M|G|\infty$ system. *International Journal of Academic Research*, 4(3), part A, 24-33.
- [21] FERREIRA, M. A. M., ANDRADE, M. (2012c). A methodological note on the study of queuing networks. *Journal of Mathematics and Technology*, 3(1), 4-6.
- [22] FERREIRA, M. A. M., ANDRADE, M., FILIPE, J. A (2012). The age or excess of the $M/G/\infty$ queue busy cycle mean value. *Computer and Information Science*, 5(5), 93-97. <http://dx.doi.org/10.5539/cis.v5n5p93>
- [23] FERREIRA, M. A. M. (2013). A $M|G|\infty$ queue busy period distribution characterizing parameter. *Computer and Information Science*, 6 (1), 83-88. <http://dx.doi.org/10.5539/cis.v6n1p83>
- [24] FERREIRA, M. A. M. (2013a). The modified peakedness as a $M/G/\infty$ busy cycle distribution characterizing parameter. *International Journal of Academic Research*, 5(2), part A, 5-8.
- [25] FERREIRA, M. A. M. (2016). Results and applications in statistical queuing theory. *APLIMAT 2016 - 15th Conference on Applied Mathematics. Proceedings*. Bratislava, Slovakia, 362-375.
- [26] FERREIRA, M. A. M. (2017). A particular collection of service time distributions parameters study and impact in some $M|G|\infty$ system busy period and busy cycle parameters. *Acta Scientiae et Intellectus*, 3(1), 31-39.
- [27] FERREIRA, M. A. M., FILIPE, J. A. (2017). In the search for the infinite servers queue with Poisson arrivals busy period distribution exponential behaviour, *Int. J. Business and Systems Research*, 11(4), 453-467.
- [28] FIGUEIRA, J., FERREIRA, M. A. M. (1999). Representation of a pensions fund by a stochastic network with two nodes: an exercise. *Portuguese Revue of Financial Markets*, 1(3).
- [29] HERSHEY, J. C., WEISS, E. N., MORRIS, A. C. (1981). A stochastic service network model with application to hospital facilities. *Operations Research*, 29(1), 1-22, 1981.
- [30] KENDALL, M. G., STUART, A. (1979). *The advanced theory of statistics. Distributions theory*. London, Charles Griffin and Co., Ltd. 4th Edition.
- [31] KLEINROCK, L. (1985). *Queueing systems*. Vol. I and Vol. II. Wiley- New York.

- [32] RAMALHOTO, M. F., FERREIRA, M. A. M. (1996). Some further properties of the busy period of an $M|G|\infty$ queue. *Central European Journal of Operations Research and Economics*, 4(4), 251-278.
- [33] ROSS, S. (1983). *Stochastic Processes*. Wiley, New York.
- [34] STADJE, W. (1985). The busy period of the queueing system $M|G|\infty$. *Journal of Applied Probability*, 22, 697-704.
- [35] SYSKI, R. (1960). *Introduction to congestion theory in telephone systems*. Oliver and Boyd, London.
- [36] SYSKI, R. (1986). *Introduction to congestion theory in telephone systems*. North Holland, Amsterdam
- [37] TAKÁCS, L. (1962). *An introduction to queueing theory*. Oxford University Press, New York.
- [38] WHITT, W. (1984). On approximations for queues, I: extremal distributions. *A T & T Bell Laboratories Technical Journal*, 63(1), 115-138.

Current address

Ferreira Manuel Alberto M., Professor Catedrático
INSTITUTO UNIVERSITÁRIO DE LISBOA (ISCTE-IUL)
BRU – IUL, ISTAR-IUL
Av. das Forças Armadas, 1649-026 Lisboa, Portugal
Tel.: + 351 21 790 37 03. Fax: + 351 21 790 39 41
E-mail: manuel.ferreira@iscte.pt

Filipe José António, Professor Auxiliar
INSTITUTO UNIVERSITÁRIO DE LISBOA (ISCTE-IUL)
BRU – IUL, ISTAR-IUL
Av. das Forças Armadas, 1649-026 Lisboa, Portugal
Tel.: + 351 21 790 34 08, Fax: + 351 21 790 39 41
E-mail: jose.filipe@iscte.pt